



Full Length Research Article

Advancements in Life Sciences – International Quarterly Journal of Biological Sciences

ARTICLE INFO

Open Access



Date Received:
06/08/2023;
Date Revised:
01/02/2024;
Available Online:
18/04/2024;

Author's Affiliation:

Department of Biology, College
of Science for Women,
University of Babylon - Iraq

*Corresponding Author:

Hussein J. Hussein
Email:
huss2010feb@yahoo.com

How to Cite:

Al-Nafie FS, Hussein HJ, Al-
Rubaye AF (2024). Antifungal
efficacy of the crude alkaloid,
flavonoid, and terpenoid of
Saussurea costus (Falc.)
Lipschitz roots against
Aspergillus species isolated
from rice seeds. Adv. Life Sci.
11(2): 392-397.

Keywords:

Antifungal activity;
Saussurea costus; Alkaloids;
Flavonoids; Terpenoids

Editorial Note:

You are viewing the latest
version of this article having
minor corrections related to
the use of English language.

Antifungal efficacy of the crude alkaloid, flavonoid, and terpenoid of *Saussurea costus* (Falc.) Lipschitz roots against *Aspergillus* species isolated from rice seeds

Fatima S. Al-Nafie, Hussein J. Hussein*, Abeer Fauzi Al-Rubaye

Abstract

Background: Rice is an essential component of the diets of people all over the world, and it accounts for a sizeable amount of the calories that they consume. The world's food supply suffers huge economic losses due to fungal deterioration of rice seeds. The medicinal plant *Saussurea costus* contains bioactive chemicals and is well-known for its many therapeutic uses. There are a number of species of *Aspergillus* that are considered pathogens of vegetables and fruits. One of these species is *Aspergillus niger*. The purpose of this current investigation was to investigate the impact of the crude alkaloid, flavonoid, and terpenoid chemicals derived from *Saussurea costus*' roots on *Aspergillus* species isolated from rice seeds obtained from various markets in Hillah, Iraq.

Methods: In vitro, antifungal activity against *Aspergillus* species was achieved using the food poisoning method. Three concentrations 5, 10, and 15mg/ml of each crude compound were prepared and compared to a positive control represented by Carbendazim 500g/l and a negative control represented by 10% dimethyl sulfoxide.

Results: The objective of this study was to manage the *Aspergillus* species growth detected in rice seeds by utilizing secondary compounds derived from the roots of *Saussurea costus*. The study's data indicated that the extracts represented by active compounds such as Alkaloids, Flavonoids, and Terpenoids obtained from *Saussurea costus* roots exhibited a significant decrease in the of *Aspergillus* species growth, particularly at a concentration of 100mg/ml, in contrast to the negative control. This effect was equivalent to that of the positive control, Carbendazim 500g/l.

Conclusion: Finally, *Saussurea costus* roots have shown the most efficacy in controlling *Aspergillus* species, particularly through the presence of Alkaloid, Flavonoid, and Terpenoid chemicals at a concentration of 100mg/ml.



Introduction

Oryza sativa L., represents over 50% of an essential dietary constituent of the people around the world. It contributes approximately 55 to over 79% of the entire caloric eating for individuals residing in South and Southeast Asia, and Latin America, in other places, it is regarded as a valuable cash yield [1]. Rice is crucial for ensuring food security in Iraq and various other countries [2]. Molds that ruin food and feed result in significant global economic losses. Approximately 5 to 10% of the global food supply is lost as a result of fungal decay [3]. Over 100 distinct fungal species have been found on rice seeds thus far. However, the degree of these effects varies based on the timing of sample, location, and different types [4]. *Saussurea costus* (Falc.) Lipschitz, is a species of the family *Asteraceae* and is a species native to the Himalayas, specifically found at heights ranging from 2700 to 4000 meters in Lahul Valley of Himachal Pradesh, Garhwal of Uttarakhand and Kashmir, and has another name *Saussurea lappa* C.B. Clarke [5, 6]. *S. costus* is a renowned and significant medicinal plant extensively utilized in diverse indigenous medical systems to treat a range of conditions, including asthma, inflammatory illnesses, ulcers, and stomach problems. The primary phytoconstituents of this plant are sesquiterpene lactones. Various pharmacological experiments conducted in both laboratory and animal models have provided strong evidence of *S. costus*' capacity to display anti-inflammatory, antiulcer, anticancer, and hepatoprotective properties. These findings support the traditional applications of this plant. The plant contains dehydrocostus lactone, cynaropicrin, and costunolide which have been identified as possible bioactive compounds. Given the notable biological action of *S. costus* and its ingredients, it is fitting to explore their potential as a pharmaceutical treatment [7]. Fresh roots obtained from *S. costus* plant contain lactone, lappadilactone, dehydrocostus, cynaropicrin, and germacrenes compounds [8]. *Aspergillus* is among the top three fungal genera that have a substantial effect on food deterioration and the production of mycotoxins. The other two key genera are *Fusarium* and *Penicillium*. Among these, *Aspergillus niger* is the sole public species that poses a solemn threat as a pathogen in some vegetables and fruits [9]. Instances of mycotoxin production by *A. niger* strains are infrequent [10]. *Aspergillus* species are capable of producing significant mycotoxins, specifically aflatoxins which are generated by *A. flavus* and similar species, and ochratoxin A, which is formed by *A. ochraceus* and associated species, as well as *A. carbonarius* [11]. The objective of this investigation was to manage the growth of *Aspergillus* spp. found in Rice seeds by

employing secondary metabolites derived from the roots of *S. costus*.

Methods

Plant active part: Roots belonging to *S. costus* had been procured from indigenous markets, and thereafter recognized based on their characteristics [12], The roots of this plant were cleansed, desiccated, and stored under [13], (Fig: 1).



Figure 1: *Saussurea costus* (Falc.) Lipschitz roots.

The crude alkaloid components was extracted according to the method reported in [14].

The crude flavonoid components was extracted according to the method reported in [15].

The Crude Terpenoid components was extracted according to the method reported in [16]. A 100 mg/ml stock solution of Alkaloid, Flavonoid, and Terpenoid was made in a 10% Dimethyl Sulfoxide (DMSO) solution. The solution was then sterilized using a Millipore filter with a pore size of 0.22 μ m and stored at -20°C until it was ready for use [17].

Rice seed collection: Rice seeds were collected from various parts of local shops in the area of Babil, namely in Hillah-Iraq, 2022.

Isolation and diagnosis: To separate the *Aspergillus* fungi, a total of one-hundred seeds were randomly selected from each sample. The sterilization process involved treating it with a 2% solution of sodium chlorate for 2 minutes. Subsequently, it was rinsed

twice with sterile distilled water to eliminate any remaining sterile substances. Finally, it was dried using sterile filter paper. The specimen was carefully transferred using sterilized forceps onto 9cm Petri dishes that contained 20 ml of preprepared Dextrose Agar of Potato (PDA) supplemented with 50mg/l of chloramphenicol to inhibit bacterial growing [18], each dish was seeded with 5 seeds, and each sample was replicated three times. The dishes were incubated at 25°C for 5 to 7days. The fungi that were present in the seeds were subsequently isolated using secondary cultures in order to be identified. The isolated were then identified using the keys in [19, 20]. The isolates were stored in aseptic glass containers filled with Nutrient Agar media. The containers' glass had been kept at 25°C for a duration of 7days, thereafter deposited in a fridge-freezer 4°C until they were used.

Antifungal assay: The PDA medium was prepared and sterilized. Then, a volume of (1, 2, 3ml) of each plant extract was placed in the center of the Petri dishes. The volume had been completed to 20ml with PDA medium to achieve the desired final volume concentrations of 5, 10, and 15 mg/ml of the secondary metabolites under study. Later the medium hardened entirely, a disc about of 5mm in diameter of a 7-day-old of the test fungus was placed in the center of the Petri plates under sterile conditions. The plates were then incubated at a temperature of 25±2C° for 7days. Simultaneously, 0. 02ml of an antibiotic solution had been added to each assay plate to check for bacterial contamination, as recommended by the suggested protocol [21]. The positive control utilized in the experiment was Carbendazim, a fungicide with a concentration of 500g/l. The negative control, on the other hand, was dimethyl sulfoxide with a concentration of 10%. Data was documented on the 7th day. The measurement of the colony's diameter had been expressed in millimeters. A control had been used as a medium without extract. Three replicates had been maintained for each treatment. The toxicity of the f extracts had been determined by calculating the percentage of inhibition in mycelial growth, by means of the specified formulation [22].

$$\text{Percent Inhibition\%} = [\text{dc} - \text{dt} / \text{dc}] \times 10^2$$

dc=The mean increase in colony mycelial diameter under normal conditions.

dt=The mean increase in colony mycelial diameter within the experimental treatment.

Statistical analysis: The data for treatments had been obtained from 3 replicates. The data underwent an analysis of variance by means of the SPSS, 20.0 version. A design of completely randomized had been

employed, the least significant difference (L.S.D.), test had been conducted at a significance level of Probability ≤ 0.05.

Results

Alkaloid antifungal crude compounds extracted from *S. costus* roots in contradiction of *Aspergillus* spp. isolated from seeds of Rice had been assessed in Table, 1. The findings demonstrated that Alkaloid secondary metabolites derived from *S. costus* roots exhibited a significant decline at P≤ 0.05 of *Aspergillus* spp. growth and the same effect as the positive control. Antifungal action was applied at (5.0, 10.0, and 15.0) mg/ml. mycelial reduction ranging from 66.0% in 5.0 mg/ml, 71.6% in 10.0 mg/ ml, and 100% in 15.0 mg/ml, (Figure 2: A, B), compared with negative control treatment represented by 10% DMSO and positive fungicide Carbendazim 500g/l control (Figure 2: C) where reduction percentage was 0.00% for control treatment and 100% for fungicide Carbendazim 500g/l positive control treatment, (Figure 2: D).

Concentrations	Alkaloid Antifungal Reduction%
N. Control	0.0±0.00
5.0mg/ml	66.0±1.00
10.0mg/ml	71.0.6±0.57
15.0mg/ml	100.0±0.00
P. Control	100.0± 0.00
L.S. D.	0.939

Table 1: Alkaloid Antifungal of *S. Costus* root against *Aspergillus* species.

Similarly, the crude Flavonoid compounds exhibited a 62.3% suppression of mycelial growth at a concentration of 5.0 mg/ml, and complete inhibition at concentrations of 10.0 mg/ml and 15.0 mg/ml, as indicated in table 2. Therefore, it exhibited substantial differences in comparison to the control treatment, as well as producing a similar effect to the positive control, (Figure 2: E, F).

Concentrations	Flavonoid Antifungal Reduction %
N. Control	0.0±0.00
5.0 mg/ml	62.3±0.57
10.0 mg/ml	100.0±0.00
15.0 mg/ml	100.0±0.00
P. Control	100.0±0.00
L.S. D.	0.470

Table 2: Activity of Flavonoid Antifungal of *S. Costus* root against *Aspergillus* species.

The crude Terpenoid compounds showed notable efficacy at three doses (5.0, 10.0, and 15.0 mg/ml), in comparison to the negative control. Additionally, they demonstrated a similar impact as the positive control at concentrations of 10.0 and 15.0 mg/ml, (table 3). The inhibitory action exhibited a percentage of 34.1% at a concentration of 5.0 mg/ml, while reaching 100% at concentrations of 10.0 and 15.0 mg/ml, respectively, (Figure 2: G, H). Finally, secondary metabolites,

including Alkaloid products, Flavonoid products, and Terpenoid products, isolated from *S. Costus* roots exhibit similar effects to the positive fungicide Carbendazim 500g/l control.

Concentrations	Terpenoid Antifungal Reduction %
N. Control	0±0.00
5.0mg/ml	34.1±1.04
10.0mg/ml	100.0±1.00
15.0mg/ml	100.0±0.00
P. Control	100.0±0.00
L.S. D.	1.174

Table 3: Terpenoid Antifungal *S. Costus* root against *Aspergillus* species.




Figure 2: Effect of antifungal efficacy in growth of *Aspergillus* spp. A- Alkaloid antifungal at 10.0mg/ml; B- Alkaloid antifungal at 15.0mg/ml; C- growth of *Aspergillus* spp. in the control treatment; D- growth of *Aspergillus* spp. in the fungicide Carbendazim 500g/l treatment; E- Flavonoid antifungal at 10.0mg/ml; F- Flavonoid antifungal at 15.0mg/ml; G- Terpenoid antifungal at 10.0mg/ml; H- Terpenoid antifungal at 15.0mg/ml.

Discussion

Medical plants are regarded as a crucial resource for the treatment and prevention of numerous types of diseases [23]. Every plant has numerous essential components that have potential use in the field of medicine and can contribute to the creation of various pharmaceuticals [24]. The current investigation demonstrated that chemical secondary compounds, including Alkaloids, Flavonoids, and Terpenoids, derived from *S. costus* roots exhibited anti-fungal properties counter to *Aspergillus* spp. obtained from rice seeds collected from several local markets of Hillah city located in Iraq. Chemical secondary compounds derived from several active parts of different medicinal plants such as *Lactuca serriola* leaves, *Lepidium sativum* leaves, *Myrtus Communis* leaves, *Cassia senna*

leaves, *Ricinus communis* leaves, *Cassia didymobotrya* leaves, *Melia azedarach* leaves, *Dianthus caryophyllus* flower buds, and *Salvia hispanica* seeds, exhibit antibacterial properties capable of suppressing various pathogenic microorganisms obtained from diverse clinical samples [25-33]. Hussein *et. al.* [34] it has been shown that phytochemical substances isolated from the unicellular primitive plant *Chlorella vulgaris* have the capacity to counteract harmful bacteria by acting as antibacterial agents. Kamal *et. al.* [35] Used *Hibiscus sabdarifa* extracts for controlling pathogenic microorganisms such as *E. coli* and *Proteus* sp. Kamal *et. al.* [36] Employed *Ficus carica* for inhibiting *Pseudomonas aeruginosa* and *Escherichia coli* pathogenic bacteria. AL-Masoodi *et. al.* [37] Utilized substances derived from *Curcuma longa* and *Boswellia carteri* to manage *Fusarium* spp. obtained from corn seeds. Referring to [38, 39] Employed terpenoids derived from seeds of *Carthamin tinctorius* and flavonoids obtained from *M. Communis* leaves to combat *Aspergillus* spp. identified from preserved seeds of medicinal plant. Secondary compounds found in *M. Communis* leaves, including alkaloids and flavonoid compounds, are considered valuable for their ability to inhibit pathogenic microorganisms that are isolated from hemodialysis fluid specimens [40]. Radhi *et. al.* [41] Employed extracts from *Callistemon viminalis* leaves to manage isolates of Urinary Tract Infections. *Allium sativum* is considered a reliable means of suppressing some bacterial species that have been isolated from people infected with the Corona Virus [42]. The flower buds of *D. caryophyllus* contain secondary metabolite chemicals, specifically terpenoids and flavonoids, which have potent antifungal properties against *Candida* species [43]. *S. costus* was regarded as a therapeutically significant plant, the numerous compounds extracted from it have remedial properties [12]. The roots of *S. costus* contain alkaloids and terpenoids that exhibit potent antifungal properties against *Candida* species [44]. Several chemicals were extracted from the roots of *S. costus* and evaluated against four species of *Aspergillus* such as *A. niger*, *A. ochraceus*, *A. versicolor*, *A. flavus*, and two species of *Penicillium* such as *P. ochrochloron*, *P. funiculosum*, and other species such as *Cladosporium cladosporioides*, *Alternaria*, and *Trichoderma viride*. The chemical exhibited anti-fungal properties ranging from temperate to strong [45]. The polyphenol content of *S. costus* was found to be highest in the ethanol and ethyl acetate extracts, n-butanol extract came next, while n-hexane extract came last, these compounds exhibit antibacterial and antifungal properties against a wide range of microorganisms [46]. *Saussurea lappa* had the most potent antifungal action against *A. flavus*, with *Trapa natans* and *Mangifera indica* showing lesser

efficacy [47]. The methanolic extract of *S. costus* roots is abundant in many components including flavonoids, terpenoids, alkaloids, phenols, polyphenols, tannins, resins, coumarins, quinines, steroids, and cardiac glycosides exhibited antifungal properties against *A. niger* strain (ATCC6275), [48]. Conversely, the mechanism by which Alkaloids exert their antifungal effects is typically pleiotropic, involving inhibition of protein synthesis and intercalation into fungal DNA, or by improving the production of fungal inhibitors. [49]. Terpenoids decreased the number of mitochondria, hence altering the levels of production of ATP and reactive oxygen species. Additionally, it has been found that triterpenoids have greater antifungal action in comparison to tetraterpenoids [50]. Secondary products such as Terpenoid and flavonoid exert their own effects by destabilizing microbial membrane components [51]. Flavonoids commonly hinder the growth of fungi by diverse ways, such as disrupting the plasma membrane, causing dysfunction in the mitochondria, and inhibiting processes including cell wall production, cell division, RNA, and protein synthesis, as well as the efflux-mediated pumping system. [52]. Therapeutic herbs exhibit anti-fungal properties through numerous methods. They disrupt the cell membrane, leading to an injury of membrane integrity. They also hinder DNA transcription and decrease the number of fungal cells. Additionally, they block the action of fungal antioxidant enzymes and prevent the creation of fungal biofilms [53, 54]. Finally, the anti-fungal activity of *S. costus* roots may be attributed to alkaloids, flavonoids, and terpenoids. These compounds could affect proteins and DNA synthesis, as well as impair membrane permeability and disturb metabolic activity.

The roots of *S. costus* include alkaloids, flavonoids, and terpenoids that exhibit potent antifungal properties against *Aspergillus* spp. that exist in rice seeds.

Author Contributions

Fatima S. Al-Nafie: study design, Hussein J. Hussein: Refinement of study design and supervision, Abeer Fawzi Al-Rubaye: Refinement of study design and supervision.

Conflict of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

References

- Gleick PH. Water in crisis. New York: Oxford University Press; a guide to the world's freshwater resources. Pacific Institute for Studies in Development, Environment, and Security. Stockholm Environment Institute, Oxford University Press, (1993); New York (USA); 13-24.
- Khush GS. What it will take to Feed 5.0 Billion Rice consumers in 2030. Plant. Molecular Biology, (2005); 59: 1–6.
- Pitt JI, Hocking AD. Fungi and food spoilage. Chapman and Hall, New York, USA. (1997); 593.
- Javaid MS, Wahid A, Idrees M, Gill MA, Saleem A. Seed mycoflora studies in rice. Pakistan Journal Phytopathology, (2002); 14: 132-134.
- Kaul MK. Medicinal plants of Kashmir and Ladakh: temperate and cold arid Himalaya. Indus publishing, (1997); 1-173.
- Nadkarni KM. Indian materia medica, popular Book Depot. Bombay. 1954; 7:946-8.
- Pandey MM, Rastogi S, Rawat AK. *Saussurea costus*: botanical, chemical and pharmacological review of an ayurvedic medicinal plant. Journal of Ethnopharmacology, (2007); 110(3):379-90.
- Taniguchi M, Kataoka T, Suzuki H, Uramoto M, Ando M, et al. Costunolide and dehydrocostus lactone as inhibitors of killing function of cytotoxic T lymphocytes. Bioscience, biotechnology, and biochemistry, (1995); 59(11):2064-7.
- Pitt JI, Hocking AD. Fungi and Food Spoilage. New York: Elsevier, (2009); 338.
- Cabañes FJ, Bragulat MR. Black aspergilla and ochratoxin A-producing species in foods. Current Opinion in Food Science, (2018); 23:1-10.
- Taniwaki MH, Pitt JI, Magan N. *Aspergillus* species and mycotoxins: occurrence and importance in major food commodities. Current Opinion in Food Science, (2018); (1): 23:38-43.
- Kamalpreet LK, Singh A, Kaur J, Kaur N. A brief review of remedial uses of *Saussurea lappa*. Journal of Pharmacognosy and Phytochemistry, (2019); 8(3):4423-30.
- Harborne JB, Mabry TY, Marby H. Physiology and function of flavonoids. Academic Press, New York. (1975); 970.
- Harborne JB. Phytochemical methods, London. Chapman and Hall, Ltd. (1973); 49-188.
- Boham BA, Kocipai-Abyazan R. Flavonoids and condensed tannins from leaves of Hawaiian *Vaccinium vaticulatum* and *V. calycinum*. Pacific Science, (1974); 48:458-463.
- Harborne JB. Phytochemical methods. Chapman and Hall, New York. 2nd ed. (1984); 288.
- Al-Jassani MJ. *Tropaeolum Majus* Leaves Extract as an Antifungal, Antiaflatoxigenic and Antiaflatoxin Agent. Journal of Global Pharma Technology, (2017); 12(09):328-333.
- Pitt JI, Hocking AD. Fungi and Food Spoilage. 3rd Edn. Blackie Academic Professional, London (1997); 45.
- Barnett HL, Hunter BB. Illustrated genera of imperfect fungi. Burgess. publ. Co. Minnesota, 3rd Edn. (1972); 241.
- Domsch KH, Gams W, Anderson TH. Compendium of soil fungi. Academic press. London, New York, Toronto, Sydney, San Francisco, (1980); Vol. 1; 570.
- Gupta S, Banerjee AB. A rapid method of screening antifungal antibiotic producing plants. Indian journal of experimental biology, (1970); 8(2):148-9.
- Singh J, Tripathi NN. Inhibition of storage fungi of black gram (*Vigna mungo* L.) by some essential oils. Flavour and Fragrance Journal, (1999); 14(1):1-4.
- Rakotoarivelo NH, Rakotoarivony F, Ramarosandratana AV, Jeannoda VH, Kuhlman AR, et al. Medicinal plants used to treat the most frequent diseases encountered in Ambalabe rural community, Eastern Madagascar. Journal Ethnobiology Ethnomedicine, (2015); 11(1):68.
- Yuan H, Ma Q, Ye L, Piao G. The traditional medicine and modern medicine from natural products. Molecules, (2016); 21(5): 559
- Al-Marzoqi AH, Hussein HJ, Al-Khafaji NM. Antibacterial Activity of the Crude Phenolic, Alkaloid and Terpenoid Compounds Extracts of *Lactuca serriola* L. on Human

- Pathogenic Bacteria. Chemistry and Materials Research, (2015); 7(1): 8-10.
26. Al-Marzoqi AH, Al-Khafaji NM, Hussein HJ. In vitro Antibacterial Activity Assessment of the crude Phenolic, Alkaloid and Terpenoid compounds extracts of *Lepidium sativum* L. on Human Pathogenic Bacteria. International Journal of ChemTech Research, (2016); (4): 529-32.
 27. Hussein HJ, Al-Khafaji NM, Al-Mamoori AH, Juaifer WA, Al-Marzoqi AH, et al. Antimicrobial Effect of the Crude Phenolic, Alkaloid and Terpenoid Compounds Extracts of *Myrtus Communis* L. against Human Gram-Negative Pathogenic Bacteria. Journal of Global Pharma Technology, (2017); (8): 130-3.
 28. Hussein HJ, Al-Khafaji NM, Al-Mamoori AH, Al-Marzoqi AH. Evaluation of in vitro antibacterial properties of the crude Phenolic, Alkaloid and Terpenoid extracts of *Cassia senna* L. against Human gram-negative Pathogenic Bacteria. Plant archives, (2018); 18(1): 354-6.
 29. Hussein HJ, Kaizal AF, Al-Khafaji NM, Sadiq ZF, Shahad AS. Evaluation of antibacterial potential of the crude Phenolic, Alkaloid and Terpenoid extracts of *Ricinus communis* L. against gram-negative Pathogenic Bacteria. Journal of Global Pharma Technology, (2018); (05): 384-388.
 30. Hussein HJ, Sahi NM, Saad AM, Altameme HJ. The Antibacterial Effect of bioactive compounds extracted from *Cassia didymobotrya* (Fresenius) Irwin & Barneby against Some Pathogenic Bacteria. Annals of Tropical Medicine and Public Health, (2019); 22(1): SPe 116.
 31. Hussein HJ, Al-Marzoqi AH. The Antibacterial efficacy of the secondary metabolites extracted from (*Melia azedarach* L.) leaves against pathogenic microorganisms isolated from burns and gingivitis infections. EurAsian Journal of Biosciences, (2020); 14(1): 561-5.
 32. Kamil SS, Hussein HJ, Al-Marzoqi AH. Evolution of Antibacterial efficacy of *Dianthus caryophyllus* L. extracts against some hospitals pathogenic bacteria. International Journal of Pharmaceutical Research, (2020); 12(3): 1274-9.
 33. Hussein HJ, Kamal SA, Sahi NM. Antibacterial Efficacy of The Seed Extract of *Salvia Hispanica* L. Against Pathogenic Bacteria Isolated from Diarrhea Cases. Biochemical and cellular archives, (2020); 20(supplement 2): 3491-94.
 34. Hussein HJ, Naji SS, Al-Khafaji NM. Antibacterial properties of the *Chlorella vulgaris* isolated from polluted water in Iraq. Journal of Pharmaceutical Sciences and Research, (2018); 10(10): 2457-60.
 35. Kamal AS, Hussein HJ, Tolaifeh ZA. Antibacterial potential of *Hibiscus sabdarifa* L. against some Enterobacteriaceae: in vitro. Biochemical and cellular archives, (2019); 19 (2): 4291-4294.
 36. Kamal SA, Al-Kaim HW, Hussein HJ. Antibacterial activity of phytochemical compounds extracted from *Ficus carica* Linn. Leaves against human pathogenic bacteria. EurAsian Journal of BioSciences, (2020); 14(1):2293-8.
 37. AL-Masoodi H, Hussein HJ, Al-Rubaye AF. Antifungal activity of the two medicinal plants (*Curcuma longa* L. and *Boswellia carteri* Birdwood) against *Fusarium* species isolated from maize seeds. International Journal of Pharmaceutical Research, (2020); 12(3): 408-14.
 38. Hussain AY, Hussein HJ, Al-Rubaye AF. Antifungal Activity of the Secondary Metabolites Extracted from *Carthamus tinctorius* L. against *Aspergillus* Species Isolated from Stored Medicinal Plants Seeds in the Iraqi Markets. Clinical Schizophrenia and Related Psychoses, (2021); 15S: Doi:10.3371/CSRP.HAHH.081221.
 39. Hussain AY, Hussein HJ, Al-Rubaye AF. Antifungal Efficacy of the crude Flavonoid, Terpenoid, and Alkaloid Extracted from *Myrtus communis* L. against *Aspergillus* species isolated from Stored Medicinal plants seeds in the Iraqi Markets. Journal of Biotechnology Research Center, (2021); 15 (2):73-80.
 40. Karem MH, Al-Abedy AN, Kadhim JH. Effect of Tomato yellow leaf curl virus (TYLCV) on the content of some mineral elements in the fruits of some tomato (*Solanum lycopersicom* L.) genotypes. In IOP Conference Series: Earth and Environmental Science, (2023); 1259(1): 012093. IOP Publishing.
 41. Radhi Safa Hasan, Kamal SA, Mohammed Sahi N, Hussein HJ. Assessment of Antibacterial Efficacy of *Callistemon viminalis* (Sol. ex Gaertn.) G. Don against Some Isolates Obtained from Urinary Tract Infections. Archives of Razi Institute, (2022); 77(2): 891-7.
 42. Hussein HJ, Kamal SA, Sahi NM, Radhi SH. Antibacterial Activity of *Allium Sativum* L. Aqueous Extract against Some Pathogenic Bacteria Isolated from Patients Infected with Corona Virus. HIV Nursing, (2023); 23(3):742-5.
 43. Karim ZM, Hussein HJ, Al-Rubaye AF. Evaluation of anticandidiasis efficacy of secondary metabolites extracted from *Dianthus caryophyllus* L. flower buds. Caspian Journal of Environmental Sciences, (2023); 21(1): 143-149.
 44. Karim ZM, Hussein HJ, Al-Rubaye AF. In vitro anticandidal activity of the secondary metabolites extracted from *Saussurea Costus* (Falc.) lipschitz roots. International Journal of Health Sciences, (2022); 6(S6): 7461-7470.
 45. Rao KS, Babu GV, Ramnareddy YV. Acylated flavone glycosides from the roots of *Saussurea lappa* and their antifungal activity. Molecules, (2007); 12(3):328-44.
 46. Deabes MM, Fatah AE, Sally I, Salem SH, Naguib KM. Antimicrobial activity of bioactive compounds extract from *Saussurea costus* against food spoilage microorganisms. Egyptian Journal of Chemistry, (2021); 64(6):2833-43.
 47. Al-Shammari SQ, Al-Abedy AN, Farhood AN. Effect of Cucumber mosaic virus infection on the chemical content of some genotypes of tomato plant (*Solanum lycopersicom* L.). In IOP Conference Series: Earth and Environmental Science, (2023); 1262(3). IOP Publishing.
 48. Abdallah EM, Qureshi KA, Ali AM, Elhassan GO. Evaluation of some biological properties of *Saussurea costus* crude root extract. Bioscience Biotechnology Research Communication, (2017); 10(4):601-11.
 49. Arif T, Bhosale JT, Kumar N, Mandal TK, Bendre RS, et al. Natural products-antifungal agents derived from plants. Journal of Asian natural products research, (2009); 11(7):621-38.
 50. Haque E, Irfan S, Kamil M, Sheikh S, Hasan A, et al. Terpenoids with antifungal activity trigger mitochondrial dysfunction in *Saccharomyces cerevisiae*. Microbiology, (2016); 85(4):436-43.
 51. Okusa P.N, Stévigny C, Duez P. Medicinal Plants: A Tool to Overcome Antibiotic Resistance? In: Varela, A.; Ibañez, J. (eds.), Medicinal plants: classification, biosynthesis and pharmacology. Nova Science Publishers, Incorporated, (2009); 315.
 52. Al Aboody MS, Mickymaray S. Anti-fungal efficacy and mechanisms of flavonoids. Antibiotics, (2020); 9(2):45.
 53. Braun PC. The effects of tea polyphenols on *Candida albicans*: inhibition of biofilm formation and proteasome inactivation. Canadian Journal of Microbiology, (2009); 55 (9): 1033 – 1039.
 54. Wu T, He M, Zang X, Zhou Y, Qiu T, et al. A structure-activity relationship study of flavonoids as inhibitors of *E. coli* by membrane interaction effect. Biochimica et Biophysica Acta (BBA) – Biomembranes, (2013); 1828(11): 2751-2756.



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License. To read the copy of this license please visit: <https://creativecommons.org/licenses/by-nc/4.0/>